

Instructions For Zoom Whitening Aftercare & At Home Trays

Congratulations! You have just experienced a revolutionary tooth whitening procedure.

Dark staining substance should be avoided because this is when your teeth are most susceptible to absorbing stain. Anything that can stain a white shirt can stain your teeth. Please avoid these items for the time you are bleaching + one additional week.

- Drinks: Coffee, Tea, Red Wine, Cola
- Condiments: Ketchup, Mustard, Red Sauces, Soy Sauce, Steak Sauce
- Food: Berries, Red Meat (Steak or Hamburger)
- Toothpaste: Colored toothpaste (Would suggest using Sensodyne toothpaste)
- Mouth rinse: Colored Mouthwash, Fluoride treatment, Chlorohexidine

Take Home Whitening

3 Days after ZOOM whitening session in office, start the Whitening gels at home

- Please make sure the custom tray is completely dry. Place a small amount of gel on each of the teeth in the custom tray. Use caution not to overload the tray. Insert the tray over the teeth and gently wipe away any excess gel. Do not leave any gel on the gums because this will cause irritation to your gums.
- Wear the bleaching trays with the gel for 2 hours
- Use all the bleaching gel given. This could last from a few days to a week
- If sensitivity is uncomfortable, stop and reduce time and/or frequency of treatment. If too uncomfortable, stop all together.

Things not to do when the bleaching trays are in your mouth:

- Do not rinse your mouth, since this may dilute the bleaching agent.
- Do not swallow the whitening gel.
- Never drink, eat, or smoke while wearing your tray.

After completing the daily bleaching session, remove the tray and gently clean it with a toothbrush. Rinse it completely, and let dry thoroughly before the next session. Excess water in the tray can dilute the gel.

Post-operative sensitivity

Some patients experience sensitivity such as “zingers” or thermal sensitivity 12 to 24 hours after Zoom! Whitening and during at home whitening. Here are things you can do to help relieve this sensitivity:

- Place tooth relief gel (can order on line) on the problem area in the take home trays and seat them over the teeth. You could do this before or after the bleaching session.
- Take over the counter pain relievers such as Advil or Tylenol
- Brushing your teeth with Sensodyne toothpaste.
- Drink plenty of water